



Ulcerative Colitis Agents Therapeutic Class Review (TCR)

February 20, 2016

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, digital scanning, or via any information storage or retrieval system without the express written consent of Provider Synergies, L.L.C.

All requests for permission should be mailed to:

Magellan Rx Management
Attention: Legal Department
6950 Columbia Gateway Drive
Columbia, Maryland 21046

The materials contained herein represent the opinions of the collective authors and editors and should not be construed to be the official representation of any professional organization or group, any state Pharmacy and Therapeutics committee, any state Medicaid Agency, or any other clinical committee. This material is not intended to be relied upon as medical advice for specific medical cases and nothing contained herein should be relied upon by any patient, medical professional or layperson seeking information about a specific course of treatment for a specific medical condition. All readers of this material are responsible for independently obtaining medical advice and guidance from their own physician and/or other medical professional in regard to the best course of treatment for their specific medical condition. This publication, inclusive of all forms contained herein, is intended to be educational in nature and is intended to be used for informational purposes only. Send comments and suggestions to PSTCREditor@magellanhealth.com.

February 2016

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
© 2004-2016 Magellan Rx Management. All Rights Reserved.

MagellanRx
MANAGEMENTSM

FDA-APPROVED INDICATIONS

Drug	Manufacturer	Indication(s)	
		Treatment	Maintenance
Oral Prodrug Forms			
balsalazide (Colazal®) ¹	generic	Mild to moderately active ulcerative colitis (UC) in patients 5 years of age and older	--
balsalazide (Giazo®) ²	Salix	Mild to moderately active UC in male patients 18 years of age and older	--
olsalazine (Dipentum®) ³	Meda	--	Maintenance of remission of UC in patients intolerant of sulfasalazine
sulfasalazine (Azulfidine®, Azulfidine EN-tabs®) ^{4,5}	generic	Mild to moderately active UC Adjunctive therapy in severe UC	Maintenance of remission of UC
		Other: Enteric-coated tablets are indicated in patients with UC who cannot take uncoated sulfasalazine tablets because of gastrointestinal (GI) intolerance Treatment of rheumatoid arthritis that has not responded adequately to salicylates or other nonsteroidal anti-inflammatory agents (NSAIDs) Treatment of pediatric patients with polyarticular juvenile rheumatoid arthritis who have not responded adequately to salicylates or other NSAIDs	

FDA-Approved Indications (continued)

Drug	Manufacturer	Indication(s)	
		Treatment	Maintenance
Oral Delayed-Release Forms			
mesalamine delayed-release tablets (Asacol® HD) ^{6*}	Actavis	Moderately active UC	--
mesalamine delayed release capsules (Delzicol®) ^{7*}	Actavis	Mild to moderately active UC in patients 5 years and older	Maintenance of remission of UC in adults
mesalamine MMX delayed release tablets (Lialda®) ⁸	Shire US	Mild to moderately active UC	Maintenance of remission of UC
mesalamine extended release capsules (Pentasa®) ⁹	Shire US	Mild to moderately active UC	--
mesalamine extended-release capsules (Apriso®) ¹⁰	Salix	--	Maintenance of remission of UC in adults
Rectal Forms			
budesonide rectal foam (Uceris®) ¹¹	Salix	Mild to moderate active UC extending 40 cm from the anal verge	--
mesalamine enemas (Rowasa®) ¹²	generic	Mild to moderately active distal UC, proctosigmoiditis, or proctitis	--
mesalamine enemas sulfite-free (sfRowasa®) ¹³	Meda	Mild to moderately active distal UC, proctosigmoiditis, or proctitis	--
mesalamine suppositories (Canasa®) ¹⁴	Actavis	Active ulcerative proctitis	--
Oral Corticosteroids			
budesonide extended-release tablets (Uceris®) ¹⁵	Santarus	Mild to moderately active UC	--

* Asacol (mesalamine delayed release tablet 400 mg), approved for treatment of mild to moderately active UC in patients 5 years of age and older and maintenance of remission of UC, has been discontinued.¹⁶ Asacol and Asacol HD delayed release tablets contain dibutyl phthalate in their enteric coatings. Delzicol delayed release capsules, which are bioequivalent to Asacol, do not; however, Delzicol is not AB-rated to Asacol as they are different formulations.

OVERVIEW

Ulcerative colitis (UC) is a chronic inflammatory disease primarily affecting the colon and rectum. The disease is characterized by superficial infiltration of the bowel wall by inflammatory white cells, resulting in multiple mucosal ulcerations and crypt abscesses. The lesions are contiguous, typically extending retrograde from the rectum, involving the descending, transverse, or the entire colon. The principal goal of treatment for UC is inducing, then maintaining, remission of the disease.

UC affects approximately 500,000 persons in the United States with an incidence of 8 to 12 per 100,000 population per year.¹⁷ The disease accounts for 250 million annual physician visits; 30,000 hospitalizations; and a loss of over 1 million workdays per year. The onset of UC is most common between 15 and 40 years of age, with a second peak between 50 and 80 years of age.

The predominant symptom of UC is diarrhea, which is usually associated with blood in the stool. Bowel movements are frequent but small in volume as a result of rectal inflammation. Other symptoms include pain in the lower quadrant or rectum. Systemic features, including fever, malaise, and weight loss are more common if a greater portion of the colon is affected. Elderly patients often complain of constipation rather than diarrhea because rectal spasm prevents passage of stool. The initial attack of UC may be fulminant with bloody diarrhea, but the disease more commonly begins indolently, with non-bloody diarrhea progressing to bloody diarrhea. Ulcerative colitis can present initially with any extent of anatomic involvement ranging from disease confined to the rectum to pancolitis. Most commonly, UC follows a chronic intermittent course with long periods of quiescence interspersed with acute attacks lasting weeks to months. However, a significant percentage of patients suffer a chronic continuous course.¹⁸

Aminosalicylates remain first-line treatment options for mild to moderate active UC.¹⁹ The mesalamine agents currently are available in oral and rectal formulations. The rectal products achieve high luminal concentrations of the active component, 5-aminosalicylic acid (5-ASA, mesalamine), while minimizing adverse events due to systemic absorption. Several aminosalicylates are available and differ only in mode of distribution throughout the small intestine and colon. Second-line therapy with a course of oral steroids is indicated in patients with mild to moderate disease that does not respond to oral and rectal mesalamine agents or in patients with moderate to severe disease.²⁰

For active ulcerative proctitis, an effective and rapid-acting approach is nightly administration of mesalamine retention enemas or suppositories, often supplemented with an oral aminosalicylate. Corticosteroid enemas can also be used. Another approach to proctitis is administration of an oral aminosalicylate alone, although therapeutic response may not be evident for 3 to 4 weeks.²¹

In 2007, the American Academy of Family Physicians (AAFP) released guidelines for the diagnosis and treatment of UC.²² The guidelines state that the incidence of colon cancer is increased with UC and achieving remission is critical in order to reduce a patient's lifetime risk. According to the AAFP guidelines, 5-ASA (mesalamine) via suppository or enema is first-line for patients with proctitis or proctosigmoiditis, respectively; patients unable to tolerate rectally administered 5-ASA therapy may try oral preparations, although response times and remission rates may not be as favorable. In patients with greater colonic involvement, first-line therapy should include oral mesalamine and oral corticosteroids to maintain remission. To prevent relapse of the disease, patients with UC may be given nonpathogenic *Escherichia coli* instead of 5-ASA. Symptoms refractory to oral mesalamine or oral corticosteroids may be treated with azathioprine (Imuran®) or infliximab (Remicade®). For patients

who fail to improve with the maximal dosage of 5-ASA compounds or who cannot tolerate the side effects, oral steroid therapy may be used. Prednisone is given in dosages of 40 to 60 mg per day. With the full-dose continued until symptoms are completely controlled (usually 10 to 14 days) followed by a gradual taper. Long-term oral steroid use is not recommended for chronic maintenance due to significant side effects.

The 2010 Practice guidelines of the American College of Gastroenterology state differences in treatment are based on disease severity.²³ The recommendations for maintenance and remission in distal disease include mesalamine suppositories in patients with proctitis and mesalamine enemas in patients with distal colitis (even when dosed as infrequently as every third night). Sulfasalazine, mesalamine compounds, and balsalazide are also effective in maintaining remission. The combination of oral and rectal mesalamine is more effective than either one alone. Patients with active disease (mild to moderate extensive colitis) should begin therapy with oral sulfasalazine or an alternate aminosalicylate.

In patients with severe or refractory UC symptoms, oral corticosteroids are indicated. Corticosteroids, while highly efficacious in short-term use, have numerous adverse effects, especially in the elderly, which preclude long-term use.²⁴ Patients who respond to oral prednisone and can be fully withdrawn from the drug should be maintained on an aminosalicylate. For patients with corticosteroid-dependent or corticosteroid-refractory disease, immunosuppression with azathioprine or mercaptopurine may prevent colectomy.²⁵ Adalimumab (Humira®), golimumab (Simponi®), and infliximab (Remicade), are Tumor Necrosis Factor Inhibitors (TNF-inhibitors), approved for inducing and maintaining achieving clinical response/remission in patients with moderate to severe active UC who fail conventional therapy.^{26,27,28} Vedolizumab (Entyvio®) is an integrin receptor antagonist approved for inducing and improving clinical response/remission in patients with moderate to severe active UC who showed an inadequate response to or were intolerant of treatment with a TNF inhibitor, immunomodulator or corticosteroid.²⁹ Aminosalicylates are the focus of this review.

PHARMACOLOGY

The first oral aminosalicylate developed, sulfasalazine, consists of a sulfapyridine carrier moiety linked to 5-ASA via an azo bond.³⁰ Colonic bacteria cleave the azo bond, converting sulfasalazine into sulfapyridine and 5-ASA moieties.³¹ While the sulfapyridine is absorbed and excreted in the urine, the 5-ASA component stays in the colon and is excreted in the feces. Although the specific mechanism is unknown, the intraluminal activity of 5-ASA produces a local therapeutic effect.^{32,33} Mucosal production of arachidonic acid metabolites, through cyclooxygenase and lipoxygenase pathways, is increased in patients with chronic inflammatory bowel disease. 5-ASA may decrease inflammation by blocking production of arachidonic acid metabolites in the colon.³⁴

Subsequently oral agents were developed to enhance 5-ASA delivery to the colon and reduce the incidence of adverse events.³⁵ The formulations fall into 3 categories: azo-bonded prodrug formulations (Colazal, Giazol, Dipentum), delayed-release formulations achieved by pH shift (Apriso, Asacol HD, and Lialda) or controlled-release formulations (Pentasa). The azo-bonded prodrugs are similar to sulfasalazine, and colonic bacteria are required to cleave the azo bond and release the active 5-ASA moiety.^{36,37} Effectiveness of delayed and controlled-release formulations may be variable because release of mesalamine is pH-dependent. As a result, early release increases absorption of 5-ASA in the proximal small intestine, increasing systemic exposure to 5-ASA and possible nephrotoxicity.³⁸ Apriso capsules have the Intellicor extended release delivery technology that combines an enteric pH-dependent coating giving a delayed release starting at a pH of 6 followed by a polymer matrix core that provides for extended release.³⁹ Asacol HD tablets are coated with a pH-sensitive acrylic polymer that delays the release of 5-ASA. Lialda uses MMX technology, a pH-dependent gastro-resistant coating, to delay the release of 5-ASA from the tablet core to the colon. Pentasa uses a water gradient to release microspheres containing 5-ASA from the capsule.

In March 2012, the FDA issued a draft guidance recommending against use of 2 specific phthalates, dibutyl phthalate (DBP) and di(2-ethylhexyl) phthalate (DEHP), as excipients due to developmental and reproductive toxicants in laboratory animals and potential for being endocrine disrupting, and affecting reproductive and developmental outcomes in humans.⁴⁰ These agents are often used as plasticizers in enteric and delayed release coatings of drug products, including a mesalamine DR product (Asacol HD). The draft guidance recommended that manufacturers reformulate products containing these excipients. A final guidance was issued in December 2012.⁴¹ Mesalamine DR (Delzicol) has been formulated without DBP but substituted with dibutyl sebacate (DBS).

Mesalamine is available as suppositories (Canasa) and enemas that deliver 5-ASA directly to the site of action. For the treatment of ulcerative proctitis, mesalamine suppositories (or corticosteroid foam), which deliver drug to the rectum, are appropriate for the treatment of up to 20 cm of distal colon. Mesalamine (or corticosteroid) retention enemas, which distribute drug to the left colon, can be used for active disease involving up to 60 centimeters of distal colon.⁴² A sulfite-free formulation of mesalamine enema (sfRowasa) has been FDA-approved.⁴³

Steroids, such as budesonide ER (Uceris), may suppress autoimmune and inflammatory responses in UC. Budesonide has a high topical glucocorticosteroid activity and substantial first-pass elimination. Uceris is a delayed and extended-release tablet, which breaks down at pH \geq 7. Uceris uses MMX technology. Budesonide is available as rectal foam (Uceris rectal foam), can be used for active mild to moderate distal ulcerative colitis extending up to 40 centimeters from the anal verge.

PHARMACOKINETICS^{44,45,46,47,48,49,50,51,52,53,54,55,56,57,58}

All aminosalicylate oral products are designed to release 5-ASA for action in the intestine so systemic absorption is intended to be minimal. Absorbed 5-ASA and its metabolites are excreted in the urine. The majority of 5-ASA remains in the colonic lumen and is excreted in feces. The apparent half-life of 5-ASA can range from 2 to 15 hours due to the different formulations of the drugs.

Drug	Delivery Mechanism	Bioavailability (%)
Oral Prodrug Forms		
balsalazide (Colazal, Giazol)	Delivered to the colon intact then bacteria cleave the compound to release 5-ASA	low and variable
olsalazine (Dipentum)	Rapidly converted in the colon to molecules of 5-ASA by bacteria and the colon's low prevailing redox potential	2.4
sulfasalazine (Azulfidine, Azulfidine En-Tabs)	Metabolized by intestinal bacteria to 5-ASA and sulfapyridine; site of delivery is the colon Azulfidine En-Tabs contain a cellulose acetate phthalate coating that retards disintegration in the stomach	<15
Oral Delayed-Release Forms		
mesalamine delayed-release tablets (Asacol HD)	Acrylic-based resin coating delays 5-ASA release until tablet reaches the terminal ileum and beyond; pH dependent release at pH ≥ 7	20–25
mesalamine delayed release capsules (Delzicol) [†]	Capsules contain acrylic based resin, Eudragit S (methacrylic acid copolymer type B, NF), which delays 5-ASA release until capsule reaches the terminal ileum and beyond; pH dependent release at \geq pH 7	28
mesalamine MMX tablets (Lialda)	pH dependent polymer film breaks down at pH ≥ 6.8 , in terminal ileum where mesalamine begins to be released from tablet core; tablet core has hydrophilic and lipophilic excipients that provide for extended release of mesalamine	21–22
mesalamine extended release capsules (Pentasa)	Ethylcellulose-coated, controlled release formulation releases 5-ASA throughout the intestinal tract	20–30
mesalamine extended release capsules (Apriso)	Intellicor extended-release delivery technology that combines an enteric pH-dependent coating which provides for a delayed release starting at a pH of 6 with a polymer matrix core that enables extended release	21–44

Pharmacokinetics (continued)

Drug	Delivery Mechanism	Bioavailability (%)
Rectal Forms		
budesonide rectal foam (Uceris)	Rectal Administration	60
mesalamine enemas (Rowasa, sfRowasa)	Rectal administration	10–30
mesalamine suppositories (Canasa)	Rectal administration	variable
Oral Corticosteroids		
budesonide extended release tablets (Uceris) ⁵⁹	pH dependent enteric coated delayed release tablets with a polymer coating that dissolves at pH ≥7 with an extended release tablet core	10–20

† Delzicol 400 mg capsules are bioequivalent to Asacol 400 mg tablets. Two Delzicol 400 mg capsules have not been shown to be bioequivalent to 1 Asacol HD 800 mg tablets.

CONTRAINDICATIONS/WARNINGS^{60,61,62,63,64,65,66,67,68,69,70,71,72,73}

Aminosalicylates are contraindicated in patients with salicylate hypersensitivity. Sulfasalazine is also contraindicated in patients with sulfonamide hypersensitivity, porphyria, and intestinal or urinary obstruction. Budesonide ER (Uceris) is contraindicated in patients hypersensitive to budesonide or any excipients in Uceris.

Deaths associated with administration of sulfasalazine have been reported. Deaths occurred from hypersensitivity reactions, agranulocytosis, aplastic anemia, other blood dyscrasias, renal and liver damage, irreversible neuromuscular and central nervous system changes, and fibrosing alveolitis. Complete blood counts, as well as urinalysis with careful microscopic examination, should be done frequently in patients receiving sulfasalazine. Oligospermia and infertility have been observed in men treated with sulfasalazine; however, withdrawal of the drug appears to reverse the effects.

Sulfasalazine should be given with caution to patients with severe allergy or bronchial asthma. Serious skin reactions, including exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported. Drug rash with eosinophilia and systemic symptoms have also been reported, and early manifestations such as fever or lymphadenopathy may be present even though rash is not. If such signs or symptoms are present, the patient should be evaluated immediately and sulfasalazine should be discontinued if there is no alternative etiology. Adequate fluid intake must be maintained in order to prevent crystalluria and stone formation.

SJS and DRESS have also been reported with the mesalamine containing products Lialda and Pentasa.

Renal impairment has been reported in patients taking products that contain or are converted to mesalamine. Evaluate renal function prior to initiation of therapy and periodically thereafter. Patients with pyloric stenosis may have prolonged gastric retention of oral mesalamine and balsalazide, which could delay the release of drug in the colon. There have been reports of hepatic failure in patients with pre-existing liver disease who have been administered mesalamine. Lastly, mesalamine, balsalazide,

and olsalazine have been associated with an acute intolerance syndrome that may be difficult to distinguish from a UC flare. These symptoms may abate once the agent is discontinued.

Mesalamine enemas (Rowasa Rectal Suspension Enema) contain potassium metabisulfite, a sulfite which may cause life-threatening allergic-type reactions including anaphylaxis. Sulfite sensitivity is more frequent in asthmatic patients or atopic non-asthmatic persons. Overall prevalence of sulfite sensitivity in the general population is not known, but probably low. A sulfite-free mesalamine enema (sfRowasa) is available; it is proposed to be safe for use in patients with sulfite allergy.

Dibutyl phthalate (DBP) is an inactive ingredient in the enteric coating of Asacol HD brand (mesalamine) tablets. In animal studies at doses Asacol HD > 80 times the human dose based on body surface area, maternal DBP was associated with external and skeletal malformations and adverse effects on the male reproductive system. Asacol HD should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Chronic glucocorticosteroid use may cause hypercorticism, adrenal suppression, and can reduce the response of the hypothalamus-pituitary adrenal (HPA) axis to stress, such as surgery. Patients who are switched from glucocorticosteroids with greater systemic effects may undergo withdrawal, including acute adrenal suppression or benign intracranial hypertension. Consequently, adrenocortical function should be monitored and the dose of the high potency glucocorticosteroid should be reduced cautiously. Glucocorticoids can also suppress the immune system causing increased susceptibility to infection. Consequently, exposure to transmissible diseases should be avoided and use in patients with active or quiescent tuberculosis infection, untreated fungal, bacterial, systemic viral or parasitic infections should be done cautiously, if at all. Liver dysfunction may decrease elimination and increase bioavailability resulting in increased toxicity. Caution should be observed in patients with hypertension, diabetes, osteoporosis, peptic ulcer, glaucoma or cataracts, or with a family history of diabetes or glaucoma, or with any other condition where glucocorticoids may have unwanted effects.

Mesalamine, a metabolite of sulfasalazine, may spuriously elevate test results for urinary normetanephrine when measured by liquid chromatography with electrochemical detection when testing for pheochromocytoma.

DRUG INTERACTIONS^{74,75,76,77,78,79,80,81,82,83,84}

Antacids: Mesalamine extended-release capsules (Apriso) depend on pH for dissolution of the coating of the granules so concomitant use with antacids should not occur.

Digoxin: Sulfasalazine, in doses more than 2 g daily, reduces the oral absorption of digoxin by 25%. It is unclear if other aminosalicylates have any significant effect on digoxin absorption.

Folic acid: Sulfasalazine can inhibit the absorption of folic acid; supplementation of folic acid may be required.

Phenytoin: Sulfasalazine can displace highly protein-bound drugs such as phenytoin.

Warfarin: Salicylates may displace warfarin from protein binding sites leading to hypoprothrombinemia. This dose-related interaction has been reported with olsalazine and sulfasalazine.

Nephrotoxic Agents, Including Non-Steroidal Anti-Inflammatory Drugs: Mesalamine use with known nephrotoxic agents, including nonsteroidal anti-inflammatory drugs (NSAIDs) may increase the risk of renal reactions.

Azathioprine or 6-mercaptopurine: Mesalamine use with azathioprine or 6-mercaptopurine may increase the risk for blood disorders.

Dissolution of the coating of budesonide ER (Uceris) is pH dependent. Consequently, drug release and absorption may be altered when budesonide ER is used with drugs that raise gastric pH (e.g., proton pump inhibitors [PPIs], histamine 2- [H2]-blockers, and antacids). Concomitant oral administration of ketoconazole (a known inhibitor of CYP3A4) caused an 8-fold increase in the systemic exposure to oral budesonide ER. If treatment with inhibitors of CYP3A4 activity (such as ketoconazole, itraconazole, ritonavir, indinavir, saquinavir, or erythromycin) is indicated, prior discontinuation of budesonide ER should be considered. Ingestion of grapefruit juice (which predominantly inhibits intestinal mucosal CYP3A4), increased systemic exposure for oral budesonide ER about 2-fold. Consequently, ingestion of grapefruit or grapefruit juice should be avoided with budesonide ER administration. Similarly, the inhibitors of CYP3A4 previously noted, may increase systemic budesonide concentrations and should be avoided with the use of budesonide rectal foam.

ADVERSE EFFECTS^{85,86,87,88,89,90,91,92,93,94,95,96,97,98,99}

Drug	Abdominal pain	Diarrhea	Fever	Headache	Nausea	Rash	Vomiting
Oral Prodrug Forms							
balsalazide (Colazal)	6–13 (3)	5–9 (3)	2–6 (0)	8–15	4–5	nr	4–10
balsalazide (Giazo)	nr	3.7 (0)	nr	nr	nr	nr	nr
olsalazine (Dipentum)	10.1 (7.2)	5.9–17 (4.8–6.7)	<1	5 (4.8)	5 (3.9)	2.3 (1.4)	1
sulfasalazine (Azulfidine)	reported	reported	less common	more common	more common	less common	more common
sulfasalazine (Azulfidine En-tabs)	8	reported	reported	9	19	13	8
Oral Delayed-Release Forms							
mesalamine delayed-release tablets (Asacol HD)	2.3	1.7	rare	4.7	2.8	reported	1.4
mesalamine MMX tablets (Lialda)	<1	<1	reported	3.4–5.6 (0.6)	nr	<1	<1
mesalamine extended release capsules (Pentasa)	1.1–1.7 (4)	3.5 (7.5)	0.9 (1.2)	2.2 (3.5)	1.8–3.1	1.3 (1.2)	1.1–1.5
mesalamine extended release capsules (Apriso)	5 (3)	8 (7)	reported	11 (8)	4 (3)	reported	nr
Rectal Forms							
budesonide rectal foam (Uceris)	nr	nr	nr	nr	2 (1)	reported	nr
mesalamine enemas (Rowasa)	8.1 (7.8)	2.1 (3.1)	3.2 (0)	6.5 (12.5)	5.8 (9.4)	2.8 (3.1)	<1
mesalamine enemas sulfite-free (sfRowasa)							
mesalamine suppositories (Canasa)	5.2	3.1	1.2 (0)	14.4	3.1	1.2 (0)	<1

Adverse Effects (continued)

Drug	Abdominal pain	Diarrhea	Fever	Headache	Nausea	Rash	Vomiting
Oral Corticosteroids							
budesonide ER tablets (Uceris) ¹⁰⁰	3.9 (1.9)	nr	nr	11.4 (10.5)	5.1 (4.3)	nr	nr

Adverse effects are reported as a percentage. Incidences reported for placebo group are shown in parentheses. Adverse effects data are obtained from package inserts and are not meant to be comparative or all inclusive. nr = not reported.

Adverse event rates with Delzicol are not available as approval was based on bioequivalence to Asacol and the data from the Asacol safety and efficacy studies.

Clinical tolerance of 3 aminosalicylate preparations [mesalamine (Asacol), olsalazine (Dipentum), and balsalazide] was assessed in a consecutive series of 43 patients with inflammatory bowel disease intolerant to sulfasalazine.¹⁰¹ Ninety-one percent of patients were able to tolerate at least 1 of the 3 preparations. Clinical tolerance of mesalamine (63%), olsalazine (70%), and balsalazide (70%) was similar. The most common adverse effects associated with the preparations were gastrointestinal in nature; diarrhea was a problem in 5 patients during treatment with olsalazine and 3 each while on mesalamine and balsalazide. Allergic reactions to aminosalicylates were uncommon; of 10 patients with rash following sulfasalazine, only 1 developed a rash with mesalamine. Results of this study indicate the vast majority of patients with inflammatory bowel disease can be managed with at least 1 of the 4 aminosalicylates, and adverse effects of sulfasalazine are multifactorial in etiology. Some adverse effects are due to the parent molecule, and some to 1 of its 2 metabolites, 5-ASA and sulfapyridine.

Renal impairment and injury including nephropathy, acute and chronic interstitial nephritis, and rarely, renal failure, have been reported in patients taking products that contain or are converted to mesalamine. In addition, exacerbation of UC symptoms has been reported upon initiation of therapy with Asacol HD as well as other mesalamine products. These symptoms usually abate once Asacol HD is discontinued. Patients with pyloric stenosis may have prolonged gastric retention of Asacol HD tablets, which could delay release of mesalamine in the colon. Lastly, there have been reports of hepatic failure in patients with pre-existing liver disease who have been administered mesalamine.

The sulfite-free mesalamine enema (sfRowasa) is proposed to cause less bowel irritation than the original Rowasa enema formulation.

In a pooled analysis of the 2 phase 3 clinical trials, there were no clinically significant differences in glucocorticoid-related adverse events between budesonide ER (Uceris) 9 mg and placebo at 8 weeks.¹⁰²

In the use of budesonide rectal foam, no clinically significant differences were reported in the overall percentage of glucocorticoid related adverse effects between budesonide foam and placebo in the course of 2 placebo-controlled trials at 6 weeks of therapy.¹⁰³ The common adverse reactions (incidence $\geq 2\%$) are decreased blood cortisol, adrenal insufficiency, and nausea.

Pediatrics

Safety and efficacy of olsalazine (Dipentum) were compared to sulfasalazine over 3 months in a multicenter, randomized, double-blind study of 56 children with mild to moderate UC.¹²⁰ Twenty-eight children received 30 mg/kg/day of olsalazine (maximum of 2 g/day) and 28 received 60 mg/kg/day of sulfasalazine (maximum of 4 g/day). After 3 months, 39% of olsalazine-treated patients were asymptomatic or clinically improved, compared to 79% of sulfasalazine-treated patients ($p=0.006$). In addition, 10 of 28 patients on olsalazine versus 1 on sulfasalazine required prednisone because of lack of response or worsening of colitis ($p=0.005$). The dose of olsalazine used in the trial was equivalent to a standard dose of sulfasalazine, but fewer patients on olsalazine improved and a greater number had progression of symptoms when compared to sulfasalazine. Adverse effects were frequent in both groups; a clinically significant difference was not detected. Safety and effectiveness of olsalazine in a pediatric population have not been established.

Sulfasalazine is approved for use in patients 6 years of age and older. While mesalamine (Asacol) was approved for the treatment of mildly to moderately active UC for 6 weeks in children 5 years of age and older with dosage based on weight, safety and efficacy of Asacol HD have not been established. Mesalamine (Delzicol) is indicated for treatment of mild to moderate active ulcerative colitis in children age 5 years and older, but is not indicated for maintenance therapy. Balsalazide 750 mg, (Colazal), is approved for use in patients 5 years of age and older, whereas balsalazide 1 gm (Giazo) is only approved in male patients 18 years and older. Colazal capsules may be opened and sprinkled on applesauce, then chewed or swallowed immediately. Other products have not been sufficiently studied in pediatric populations. Safety and effectiveness of oral budesonide ER (Uceris) as well as budesonide rectal foam (Uceris) has not been established in pediatric patients. Glucocorticosteroids, such as budesonide, may cause a reduction of growth.

Geriatrics

Reports from uncontrolled clinical studies and postmarketing reporting systems suggested a higher incidence of blood dyscrasias (e.g., neutropenia and pancytopenia) in patients who were 65 years or older who were taking mesalamine-containing products. Caution should be taken to closely monitor blood cell counts during therapy.

Clinical studies of budesonide ER (Uceris) and budesonide rectal foam (Uceris) did not include sufficient subjects aged 65 and over to determine whether they respond differently in younger subjects. In general, budesonide ER should be used cautiously in elderly patients due to the potential for decreased hepatic, renal, or cardiac function, and concomitant disease.

Pregnancy

Olsalazine (Dipentum), mesalamine (Asacol HD only), budesonide ER (Uceris), and budesonide rectal foam (Uceris) are Pregnancy Category C. Asacol HD should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Sulfasalazine (Azulfidine), balsalazide (Colazal, Giazol) and mesalamine (Delzicol) are Pregnancy Category B. There have been case reports of neural tube defects (NTDs) in infants born to mothers exposed to sulfasalazine during pregnancy. The role of sulfasalazine in these NTDs has not been established; however, oral sulfasalazine inhibits the absorption and metabolism of folic acid, which may interfere with folic acid supplementation.

Phenylketonuria (PKU)

Caution should be taken when mesalamine ER (Apriso) is administered to patients with phenylketonuria because each capsule contains aspartame equivalent to 0.56 mg of phenylalanine.

Hepatic Impairment

Patients with moderate to severe liver disease should be monitored for increased signs and/or symptoms of hypercorticism with budesonide ER (Uceris).

DOSAGES^{121,122,123,124,125,126,127,128,129,130,131,132,133,134,135}

Drug	Adults	Pediatrics	Availability
Oral Prodrug Forms			
balsalazide (Colazal)	2.25 g (three 750 mg capsules) 3 times daily for 8 to 12 weeks (Total daily dose: 6.75 g)	Children 5 to 17 yrs 2.25 g (three 750mg capsules) 3 times daily for 8 weeks OR 750 mg (1 capsule) 3 times daily for 8 weeks*	750 mg capsule
balsalazide (Giazol)	3.3 g (three 1.1 g tablets) 2 times daily with or without food for up to 8 weeks (Total daily dose: 6.6 g)	--	1.1 g tablet
olsalazine (Dipentum)	0.5 g twice daily	--	250 mg capsule
sulfasalazine (Azulfidine, Azulfidine EN-tabs)	Treatment: 3 to 4 grams daily in evenly divided doses with dosage intervals not exceeding 8 hours Maintenance: 2 grams daily	Children 6 years and older Treatment: 40 to 60 mg/kg/day divided into 3 to 6 doses Maintenance: 30 mg/kg per day divided into 4 doses	500 mg tablet 500 mg enteric coated delayed-release tablet

Dosages (continued)

Drug	Adults	Pediatrics	Availability
Oral Delayed-Release Forms			
mesalamine delayed-release tablets (Asacol HD)	1.6 g 3 times daily with or without food for 6 weeks (Total daily dose: 4.8 g)	--	800 mg delayed-release tablet
mesalamine delayed-release capsules (Delzicol)	Initial dose: 0.8 g 3 times daily for 6 weeks; 2.4 g total Maintenance dose: 1.6 g per day in divided doses for 6 months With or without food [†]	Children 12 years and older: weight-based up to a maximum of 2.4 grams/day with or without food; twice daily dosing for 6 weeks	400 mg delayed-release capsule
mesalamine MMX tablets (Lialda)	2.4 g or 4.8 g (2 to 4 tablets) once daily with a meal for up to 8 weeks [‡] Maintenance of remission: 2.4 g (2 tablets) once daily with a meal [‡]	--	1.2 g delayed-release tablet
mesalamine extended release capsules (Pentasa)	1 g 4 times a day for up to 8 weeks [§]	--	250 mg extended release capsules 500 mg extended release capsules
mesalamine extended release capsules (Apriso)	1.5 g once daily in the morning with or without food	--	0.375 g extended-release capsules

Dosages (continued)

Drug	Adults	Pediatrics	Availability
Rectal Forms			
budesonide rectal foam (Uceris)	One metered dose (2 mg) rectally twice daily for 2 weeks followed by 1 metered dose once daily for 4 weeks	--	2 aerosol canisters each containing 14 metered doses delivering 2 mg/actuation with 28 PVC applicators
mesalamine enemas (Rowasa)	4 g (60 mL) rectally at bedtime (and retained for a minimum of 8 hours) for 3 to 6 weeks	--	4 g/60 mL enema (7 and 28 unit packages) Kits include the suspension and cleansing wipes
mesalamine enemas sulfite-free (sfRowasa)	4 g (60 mL) rectally at bedtime (and retained for a minimum of 8 hours) for 3 to 6 weeks	--	4 g/60 mL enema (7 and 28 unit packages)
mesalamine suppositories (Canasa)	1 g daily at bedtime (and retained for a minimum of 1 to 3 hours) for 3 to 6 weeks	--	1,000 mg suppositories
Oral Corticosteroids			
budesonide ER tablets (Uceris) [¶]	9 mg orally once daily in the morning with or without food for up to 8 weeks	--	9 mg enteric coated delayed and extended release tablets

* Balsalazide capsules may be opened and sprinkled on applesauce; contents may be chewed.

† Mesalamine DR (Delzicol): swallow whole, do not chew, crush or break; for patients unable to swallow capsules whole, open the number of capsules required for the dose and swallow the contents whole. Delzicol has been formulated without dibutyl phthalate (DBP).

‡ Safety and efficacy of mesalamine MMX extended-release (Lialda) past 8 weeks of treatment of UC have not been established.

§ Mesalamine capsules (Pentasa): capsules may be swallowed whole or capsule may be opened and the contents sprinkled onto applesauce or yogurt. The entire contents should be consumed immediately. Capsules and contents must not be crushed or chewed.

|| The duration of mesalamine Intellicor extended-release (Apriso) use for maintaining remission of UC beyond 6 months has not been evaluated.

¶ Budesonide ER (Uceris): swallow whole, do not chew, crush or break.

CLINICAL TRIALS

Search Strategy

Articles were identified through searches performed on PubMed and review of information sent by manufacturers. Search strategy included the FDA-approved use of all drugs in this class. Randomized, controlled, comparative trials are considered the most relevant in this category. Studies included for analysis in the review were published in English, performed with human participants and randomly allocated participants to comparison groups. In addition, studies must contain clearly stated, predetermined outcome measure(s) of known or probable clinical importance, use data analysis techniques consistent with the study question and include follow-up (endpoint assessment) of at least 80% of participants entering the investigation. Despite some inherent bias found in all studies including those sponsored and/or funded by pharmaceutical manufacturers, the studies in this therapeutic class review were determined to have results or conclusions that do not suggest systematic error in their experimental study design. While the potential influence of manufacturer sponsorship/funding must be considered, the studies in this review have also been evaluated for validity and importance.

The safety and efficacy of Delzicol is based on the Asacol clinical trials.¹³⁶ Asacol has since been removed from the market, but studies including Asacol are included below as Asacol served as an active comparator for subsequent products.

mesalamine delayed-release granules (Apriso) versus placebo

Mesalamine delayed-release granules were evaluated in a double-blind, placebo-controlled trial of patients with UC in remission who took mesalamine delayed-release granules 1.5 g (n=209) or placebo (n=96) once-daily for up to 6 months.¹³⁷ The percentage of relapse-free patients at month end of treatment was higher with mesalamine than placebo (78.9% versus 58.3%; $p<0.001$) in the intent-to-treat population. Significant differences ($p\leq 0.025$) favoring mesalamine were observed for most secondary endpoints including improvement in rectal bleeding, physician's disease activity rating, stool frequency, patients classified as a treatment success, and relapse-free duration. For the mesalamine delay-release granules-treated group, 31.1% of patients withdrew from the study; UC relapse was the cause for 19.6% (n=41) of patients. For the placebo-treated group, 49% patients (n=47) withdrew from the study; 39.6% of patients (n=38) withdrew due to UC relapse. The incidence of adverse events was similar between groups. This study was sponsored by the manufacturer of Apriso, Salix Pharmaceuticals.

mesalamine delayed release (Asacol HD) 4.8 g/day versus mesalamine delayed release (Asacol) 2.4 g/day

A 6-week, multicenter, randomized, double-blind, active-control study (ASCEND III) was conducted to assess the non-inferiority of mesalamine delayed release high dose (Asacol HD) 4.8 g/day to mesalamine delayed release (Asacol) 2.4 g/day in 772 patients with moderately active UC.¹³⁸ The primary endpoint was overall improvement at Week 6 as defined by the Physician's Global Assessment (based on clinical assessments of rectal bleeding, stool frequency, and sigmoidoscopy) with no worsening in any individual clinical assessment. The primary objective of non-inferiority was met when 70% (273 of 389) of patients who received mesalamine 4.8 g/day achieved treatment success at week six compared to 66% (251 of 383) of patients receiving mesalamine 2.4 g/day. In addition, 43% of

patients receiving the higher dose of mesalamine achieved clinical remission at Week 6 compared to 35% of patients receiving the lower dose of mesalamine ($p=0.4$). A therapeutic advantage was observed for those patients who were previously treated with corticosteroids, oral mesalamine, rectal therapies, or multiple UC medications. Both regimens were well tolerated with similar adverse events.

balsalazide (Colazal) versus mesalamine delayed-release (Asacol)

A double-blind study compared the effectiveness of balsalazide and mesalamine delayed-release in the treatment of 101 patients with active moderate to severe UC.¹³⁹ Patients were randomized to receive balsalazide 6.75 g/day or mesalamine delayed-release 2.4 g/day for 12 weeks. After 2, 4, and 12 weeks, symptom control was greater in the balsalazide group. Remission rate after 12 weeks of therapy was 62% with balsalazide and 37% with mesalamine delayed-release. Median time to first day of complete relief of symptoms was 10 days for the balsalazide group and 25 days for the mesalamine delayed-release group. Adverse effects occurred in 48% of patients treated with balsalazide and 71% of those treated with mesalamine delayed-release.

A randomized, double-blind, double-dummy, parallel-group, dose-response study was performed comparing balsalazide 2.25 or 6.75 g daily and delayed-release mesalamine 2.4 g daily.¹⁴⁰ Medication was administered for 8 weeks to 154 patients with active, mild to moderate UC, the majority of who were relapsing. High-dose balsalazide was superior to low-dose in rectal bleeding, stool frequency, sigmoidoscopic score, and Physician's Global Assessment (PGA). The only significant difference observed between high-dose balsalazide and mesalamine delayed-release was more rapid onset of action as determined by a better 2-week sigmoidoscopic score for patients treated with balsalazide (55% versus 29%; $p=0.006$). Balsalazide 6.75 g daily was well tolerated, and the safety profile did not differ significantly from either balsalazide 2.25 g daily or mesalamine delayed-release 2.4 g daily.

A total of 173 patients with active, mild to moderate UC were randomized to 8 weeks of double-blind treatment with balsalazide 2.25 g or mesalamine 0.8 g, each given 3 times daily.¹⁴¹ Overall, 46% of balsalazide-treated and 44% of mesalamine-treated patients achieved symptomatic remission at endpoint. Although the median time to symptomatic remission was shorter with balsalazide (25 days) than with mesalamine (37 days), the difference was not clinically significant. Significantly more balsalazide-treated patients showed improvement in sigmoidoscopic score ($p=0.002$), stool frequency ($p=0.006$), rectal bleeding ($p=0.006$), and physician global assessment scores ($p=0.013$) by 14 days compared to mesalamine-treated patients. The difference between groups in improved sigmoidoscopic score was significant at day 28 ($p=0.002$). By day 56 and at endpoint, no significant differences between groups were detected. During the treatment period, 54% of balsalazide- and 64% of mesalamine-treated patients reported at least 1 treatment-emergent adverse event. The most common adverse events affected the gastrointestinal tract or the central and peripheral nervous systems.

The mesalamine delayed-release (Asacol) product used in the studies was manufactured and marketed by Smith Kline & French in the United Kingdom, rather than the Procter & Gamble product used in North America. Although the significance is not known, data are available from comparative *in vitro* dissolution studies to suggest slight differences exist between the 2 Asacol products.¹⁴²

balsalazide (Giazo) versus placebo

A double-blind, placebo-controlled, multicenter trial was conducted in 250 male and female adult patients with mild to moderate active ulcerative colitis.¹⁴³ Patients were randomized 2:1 to receive 8 weeks of treatment with either balsalazide 3.3 g twice daily or placebo. Disease activity was assessed using a modified Mayo Disease Activity Index (MMDAI). The primary efficacy endpoint was the proportion of patients that achieved clinical improvement and improvement in the rectal bleeding subscale of the MMDAI at the end of 8 weeks of treatment. Clinical Improvement was defined as having both greater than or equal to a 3 point improvement from baseline in the MMDAI score and greater than or equal to a 1 point improvement from baseline in the rectal bleeding sub score. At total of 55% of patients had clinical improvement with balsalazide compared to 40% on placebo, ($p=0.0237$). This difference in clinical improvement was completely due to improvement in males (57% improved with balsalazide versus 20% with placebo). There was no difference in the percentage of females with clinical improvement (54% versus 58% with balsalazide versus placebo, respectively).

olsalazine (Dipentum) versus sulfasalazine (Azulfidine)

A randomized, double-blind, 6-month study compared 3 doses of olsalazine (0.5, 1.25, and 2 g daily) and sulfasalazine 2 g daily for maintenance of remission in 162 patients with UC.¹⁴⁴ Using intention-to-treat analysis, failure rates of the different treatment groups were not significantly different (36%, 49%, and 24% for 0.5, 1.25 and 2 g olsalazine daily and 32% for 2 g sulfasalazine daily). Olsalazine and sulfasalazine showed a tendency towards lower failure rates in extended disease (28%) than in distal disease (44%). Withdrawal rate due to adverse effects was 4% with the most frequent single event being diarrhea, which occurred only in patients treated with olsalazine (2.5%, 5.2%, and 11.7% for daily olsalazine doses of 0.5, 1.25, and 2 g, respectively).

A randomized, double-blind trial compared the relapse-preventing effects of olsalazine and sulfasalazine in patients with UC over 12 months.¹⁴⁵ A total of 227 patients received either olsalazine 500 mg twice daily or sulfasalazine 1 g twice daily. A total of 197 patients completed the trial. Relapse rate after 12 months in the olsalazine group was 46.9% versus 42.4% in the sulfasalazine group (95% confidence interval (CI), -9 to 18%). Equal numbers of patients in each group withdrew from the trial because of adverse effects.

mesalamine MMX delayed-release tablets (Lialda) versus placebo

A randomized, double-blind, parallel-group, placebo-controlled trial was conducted in 280 patients with active, mild to moderate UC over 8 weeks.¹⁴⁶ Patients received mesalamine MMX delayed-release 1.2 g twice daily, 4.8 g once daily, or placebo. The primary efficacy endpoint was percentage of patients in clinical and endoscopic remission after 8 weeks of treatment. Clinical and endoscopic remission at week 8 was achieved by 34.1% and 29.2% of the mesalamine MMX delayed-release 2.4 g/day and 4.8 g/day groups, respectively, versus 12.9% of placebo patients. Mesalamine MMX delayed-release tablets given once or twice daily were well tolerated and, compared with placebo, demonstrated efficacy for induction of clinical and endoscopic remission in mild to moderately active UC.

mesalamine MMX delayed-release tablets (Lialda) versus mesalamine delayed-release tablets (Asacol)

An 8-week, double-blind, multicenter trial was conducted in 340 patients with active, mild to moderate UC comparing mesalamine MMX delayed-release 2.4 g/day or 4.8 g/day, mesalamine delayed-release 2.4 g/day given in 3 divided doses, or placebo.¹⁴⁷ The primary endpoint was proportion of patients in clinical and endoscopic remission. Remission was measured by a modified UC disease activity index of less than or equal to 1 with rectal bleeding, stool frequency scores of 0, no mucosal friability, and a greater than or equal to 1 point reduction in sigmoidoscopy score from baseline. Patients treated with mesalamine MMX delayed-release experienced significantly greater clinical and endoscopic remission rates by Week 8 versus placebo (2.4 g/day = 40.5%; 4.8 g/day = 41.2%; placebo = 22.1%). The remission rate for mesalamine delayed-release was not significantly greater than placebo (32.6%; $p=0.124$). All active treatments were well-tolerated.

A 6 month, randomized, double-blind, active-control, multicenter trial was conducted to assess the non-inferiority of once-daily mesalamine MMX (Lialda) 2.4 g/day compared with twice-daily mesalamine delayed-release (Asacol) 1.6 g/day in 826 patients with UC in maintaining endoscopic remission after 6 months.¹⁴⁸ At Month 6, 83.7% and 77.8% of patients receiving mesalamine MMX in the per-protocol and intent-to-treat (ITT) populations, respectively, maintained endoscopic remission compared with 81.5% and 76.9% of patients receiving mesalamine delayed-release (95% CI for difference -3.9%, 8.1% [for the per protocol population, PP]. Time to relapse was not significantly different between the 2 treatment groups (log-rank test, $p=0.5116$ (PP population)).

budesonide extended-release tablets (Uceris) versus placebo and budesonide

This was a randomized, double-blind, placebo-controlled study in 461 adults with active, mild to moderate UC, defined as an Ulcerative Colitis Disease Activity Index (UCDAI) of ≥ 4 and ≤ 10 and histology consistent with active UC.¹⁴⁹ Budesonide ER 9 mg and budesonide ER 6 mg (not approved in the US) were compared with, another brand of budesonide 9 mg (as reference) not approved for the treatment of UC, and compared to placebo. The primary endpoint was induction of remission after 8 weeks and remission was defined as a UCDAI score of ≤ 1 , with sub scores of 0 for rectal bleeding, stool frequency, and mucosal appearance and with a ≥ 1 point reduction in an endoscopy-only score. At the end of 8 weeks 17.4% in the budesonide ER 9 mg group, 8.3% in the budesonide ER 6 mg group, 12.6% in the budesonide 9 mg group, and 4.5% of placebo, were in remission. The difference in remission rate for budesonide ER 9 mg versus placebo was 12.9% (95% CI, 4.6 to 21.3, $p<0.025$ [$\alpha=0.025$]).

budesonide extended-release tablets (Uceris) versus placebo and mesalamine

This was a randomized, double-blind, placebo-controlled study in 509 adult patients with active, mild to moderate UC.^{150,151} Budesonide ER 9 mg and budesonide ER 6 mg (not approved in the US) was compared with mesalamine DR 2.4 g (as reference), and to placebo. The primary endpoint was induction of remission after 8 weeks. Remission was defined as a UCDAI score of ≤ 1 , with sub scores of 0 for rectal bleeding, stool frequency, and mucosal appearance and with a ≥ 1 point reduction in an endoscopy-only score. At the end of 8 weeks 17.9% of patients administered budesonide ER 9 mg, 13.2% in the budesonide ER 6 mg arm, 12.1% in the mesalamine DR 2.4 g arm, and 7.4% of placebo, were in remission. The difference in remission rate for budesonide ER 9 mg compared to placebo was

10.4% (95% CI, 2.2 to 18.7, $p < 0.025$ [$\alpha = 0.025$]). Adverse events occurred at similar rates among groups.

budesonide rectal foam (Uceris) versus placebo

Two randomized, double-blind, placebo-controlled, multicenter trials evaluated 546 adults with active mild-to-moderate distal ulcerative colitis.^{152,153} For these trials oral and rectal corticosteroids, as well as rectal 5-aminosalicylic acid (5-ASA) products were prohibited during the course of the trials, but allowed as rescue therapy. Oral 5-ASA products were allowed at doses ≤ 4.8 g/day. Patients were randomized to 267 subjects to budesonide rectal foam and the remaining 279 subjects to placebo. During each trial patients received budesonide rectal foam 2 mg or placebo twice daily for 2 weeks followed by 2 mg doses once daily for an additional 4 weeks. In each of the 2 trials, the primary endpoint was the proportion of patients who were in remission after 6 weeks of treatment with remission defined as a decrease or no change in the stool frequency subscore from baseline, a rectal bleeding subscore of 0, and an endoscopy score of 0 or 1. In each trial, a higher proportion of patients in the budesonide rectal foam group than in the placebo group were in remission at Week 6, (38.3% and 44% versus 25.8% and 22.4%, 95% CI) and had a rectal bleeding subscore of 0, (46.6% and 50% versus 28% and 28.6%, 95% CI) at Week 6.

SUMMARY

Relative tolerability and compliance must be considered in evaluation of the oral mesalamine preparations. Due to the addition of the 500 mg capsule of mesalamine controlled-release (Pentasa), daily pill burden has decreased from 16 to 8. Mesalamine controlled-release (Pentasa) is dosed 4 times a day using 8 capsules, and mesalamine delayed-release (Delzicol) is dosed 3 times a day using 6 capsules. Another formulation of mesalamine delayed-release (Asacol HD) is available at a higher strength that also allows for 3 times a day dosing using 6 tablets. One Asacol HD 800 mg tablet has not been shown to be bioequivalent to 2 Delzicol 400 mg capsules so substitution should not occur unless directed by the prescriber. Mesalamine MMX delayed-release (Lialda) is dosed once daily using 2 to 4 tablets. Mesalamine Intellicor extended-release (Apriso) is dosed once daily using 4 capsules. Mesalamine Intellicor extended-release (Apriso) and mesalamine MMX delayed-release tablets (Lialda) for maintaining remission of UC beyond 6 months have not been evaluated.

Balsalazide (Colazal, Giazol) is indicated for UC treatment. Olsalazine (Dipentum) is indicated for UC maintenance. Balsalazide (Colazal, Giazol) differs from olsalazine (Dipentum) in that balsalazide (Colazal, Giazol) appears to have a more rapid onset of effect; it may also be slightly more effective in left-sided disease. The tolerance of olsalazine (Dipentum) is often limited by a high rate of secretory diarrhea.

The adverse effect profile for sulfasalazine is less favorable than newer agents especially at higher doses. Patients with disease affecting the distal portion of the colon should use a rectal preparation either alone or in combination with oral therapy. Enemas and suppositories may provide quicker response time as well as less frequent dosing compared to oral therapy. Rectally administered mesalamine (generic, Rowasa enemas, sfRowasa enemas, Canasa suppositories) has a specific role as a non-oral treatment of distal UC, proctosigmoiditis, and proctitis. The sulfite-free mesalamine enema (sfRowasa) was FDA-approved as a formulation revision with a new trade name. It is proposed to cause

less bowel irritation and to be safe for use by patients with sulfite allergy, however, this has yet to be demonstrated clinically.

Aminosalicylates remain first-line treatment options for mild to moderate active UC. Extended-release budesonide (Uceris) and budesonide rectal foam (Uceris) offer alternatives for induction of remission in mild to moderate UC but should not be used in maintenance of remission.

REFERENCES

- 1 Colazal [package insert]. Raleigh, NC; Salix; January 2012.
- 2 Giazio [package insert]. Raleigh, NC; Salix; December 2013.
- 3 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.
- 4 Azulfidine [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.
- 5 Azulfidine EN-tabs [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.
- 6 Asacol HD [package insert]. Rockaway, NJ; Warner Chilcott; October 2013.
- 7 Delzicol [package insert]. Rockaway, NJ; Warner Chilcott; September 2015.
- 8 Lialda [package insert]. Lexington, MA; Shire US; November 2015.
- 9 Pentasa [package insert]. Lexington, MA; Shire US; October 2015.
- 10 Apriso [package insert]. Morrisville, NC; Salix; July 2009.
- 11 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.
- 12 Rowasa [package insert]. Somerset, NJ; Meda; June 2013.
- 13 sfRowasa [package insert]. Somerset, NJ; Meda; June 2013.
- 14 Canasa [package insert]. Bridgewater, NJ; Aptalis; December 2013.
- 15 Uceris [package insert]. Raleigh, NC; Salix Pharmaceuticals; January 2015.
- 16 Asacol [package insert]. Rockaway, NJ; Warner Chilcott; May 2015.
- 17 Kornbluth A, Sachar DB. Ulcerative colitis practice guidelines in adults (update): American College of Gastroenterology, Practice Parameters Committee. Am J Gastroenterol. 2010; 105:501-23. Accessed December 6, 2011.
- 18 Stenson WF. Ulcerative Colitis: Clinical Findings and Natural History. Cecil Textbook of Medicine. 2002.
- 19 Langan RC, Gotsch PB, Krafczyk MA, et al. Ulcerative colitis: diagnosis and treatment. Am Fam Phys. 2007; 76(9): 1323-1330. Available at: <http://www.aafp.org/afp/20071101/1323.html>. Accessed February 19, 2016.
- 20 Langan RC, Gotsch PB, Krafczyk MA, et al. Ulcerative colitis: diagnosis and treatment. Am Fam Phys. 2007; 76(9): 1323-1330. Available at: <http://www.aafp.org/afp/20071101/1323.html>. Accessed February 19, 2016.
- 21 Stenson WF. Ulcerative Colitis: Clinical Findings and Natural History. Cecil Textbook of Medicine. 2002.
- 22 Langan RC, Gotsch PB, Krafczyk MA, et al. Ulcerative colitis: diagnosis and treatment. Am Fam Phys. 2007; 76(9): 1323-1330. Available at: <http://www.aafp.org/afp/20071101/1323.html>. Accessed February 19, 2016.
- 23 Kornbluth A, Sachar DB. Ulcerative colitis practice guidelines in adults (update): American College of Gastroenterology, Practice Parameters Committee. Am J Gastroenterol. 2010; 105:501-23. Available at: <http://gi.org/guideline/ulcerative-colitis-in-adults/>. Accessed February 19, 2016.
- 24 Stein RB, Hanauer SB. Comparative tolerability of treatments for inflammatory bowel disease. Drug Saf. 2000; 23:429-448.
- 25 Pardi DS, Loftus EV, Camilleri M. Treatment of inflammatory bowel disease in the elderly: an update. Drugs Aging. 2002; 19:355-363.
- 26 Humira [package insert]. North Chicago, IL; AbbVie; November 2015.
- 27 Simponi [package insert]. Horsham, PA; Janssen Biotech; January 2016.
- 28 Remicade [package insert]. Horsham, PA; Janssen Biotech; October 2015.
- 29 Entyvio [package insert]. Deerfield, IL; Takeda; May 2014.
- 30 Stein RB, Hanauer SB. Comparative tolerability of treatments for inflammatory bowel disease. Drug Saf. 2000; 23:429-448.
- 31 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.
- 32 Schroder H, Campbell DES. Absorption, metabolism and excretion of salicylazosulfapyridine in man. Clin Pharmacol Ther. 1972; 13:539-551.
- 33 Kahn AK, Piris J, Trulove SC. An experiment to determine the active therapeutic moiety of sulfasalazine. Lancet. 1977; 2:892-895.
- 34 Colazal [package insert]. Raleigh, NC; Salix; January 2012.
- 35 Stein RB, Hanauer SB. Comparative tolerability of treatments for inflammatory bowel disease. Drug Saf. 2000; 23:429-448.
- 36 Chan RP, Pope DJ, Gilbert AP, et al. Studies of 2 novel sulfasalazine analogs, ipsalazide and balsalazide. Dig Dis Sci. 1983; 28:609-615.
- 37 Peppercorn MA, Goldman P. The role of intestinal bacteria in the metabolism of salicylazosulfapyridine. J Pharmacol Exp Ther. 1972; 181:555-562.
- 38 Schreiber S, Hamling J, Zehnter E, et al. Renal tubular dysfunction in patients with inflammatory bowel disease. Gut. 1997; 40:761-766.
- 39 Apriso [package insert]. Morrisville, NC; Salix; July 2009.
- 40 FDA. Draft Guidance for Industry on Limiting the Use of Certain Phthalates as Excipients in Center for Drug Evaluation and Research-Regulated Products; [Docket No. FDA-2012-D-0108]. Federal Register, Friday, March 2, 2012; 77:(42) 12852-12853. Available at: <https://www.federalregister.gov/articles/2012/03/02/2012-0699/draft-guidance-for-industry-on-limiting-the-use-of-certain-phthalates-as-excipients-in-center-for>. Accessed February 19, 2016.
- 41 FDA. Guidance for industry: Limiting the use of certain phthalates as excipients in CDER-regulated products. December 2012. <http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm330792.htm>. Accessed February 19, 2016.
- 42 Stenson WF. Ulcerative Colitis: Clinical Findings and Natural History. Cecil Textbook of Medicine. 2002.
- 43 sfRowasa [package insert]. Somerset, NJ; Meda; June 2013.
- 44 Colazal [package insert]. Raleigh, NC; Salix; January 2012.
- 45 Giazio [package insert]. Raleigh, NC; Salix; December 2013.
- 46 Delzicol [package insert]. Rockaway, NJ; Warner Chilcott; September 2015.
- 47 Pentasa [package insert]. Lexington, MA; Shire US; October 2015.
- 48 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.
- 49 Azulfidine [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.
- 50 Azulfidine EN-tabs [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.
- 51 Lialda [package insert]. Lexington, MA; Shire US; November 2015.
- 52 Apriso [package insert]. Morrisville, NC; Salix; July 2009.
- 53 Asacol HD [package insert]. Rockaway, NJ; Warner Chilcott; October 2013.
- 54 Rowasa [package insert]. Somerset, NJ; Meda; June 2013.

55 Canasa [package insert]. Birmingham, AL; Axcan Scandipharm; December 2013.

56 sfRowasa [package insert]. Somerset, NJ; Meda; June 2013.

57 Uceris [package insert]. Raleigh, NC; Salix; January 2015.

58 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.

59 Uceris [package insert]. Raleigh, NC; Salix; January 2015.

60 Colazal [package insert]. Raleigh, NC; Salix; January 2012.

61 Giazio [package insert]. Raleigh, NC; Salix; December 2013.

62 Pentasa [package insert]. Lexington, MA; Shire US; October 2015.

63 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.

64 Azulfidine [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.

65 Azulfidine EN-tabs [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.

66 Lialda [package insert]. Lexington, MA; Shire US; November 2015.

67 Apriso [package insert]. Morrisville, NC; Salix; July 2009.

68 Asacol HD [package insert]. Rockaway, NJ; Warner Chilcott; October 2013.

69 Rowasa [package insert]. Somerset, NJ; Meda; June 2013.

70 sfRowasa [package insert]. Somerset, NJ; Meda; June 2013.

71 Canasa [package insert]. Bridgewater, NJ; Aptalis; December 2013.

72 Uceris [package insert]. Raleigh, NC; Salix; January 2015.

73 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.

74 Colazal [package insert]. Raleigh, NC; Salix; January 2012.

75 Pentasa [package insert]. Lexington, MA; Shire US; October 2015.

76 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.

77 Azulfidine [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.

78 Azulfidine EN-tabs [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.

79 Lialda [package insert]. Lexington, MA; Shire US; November 2015.

80 Apriso [package insert]. Morrisville, NC; Salix; July 2009.

81 Asacol HD [package insert]. Rockaway, NJ; Warner Chilcott; October 2013.

82 Canasa [package insert]. Bridgewater, NJ; Aptalis; December 2013.

83 Uceris [package insert]. Raleigh, NC; Salix; January 2015.

84 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.

85 Colazal [package insert]. Raleigh, NC; Salix; January 2012.

86 Giazio [package insert]. Raleigh, NC; Salix; December 2013.

87 Pruitt R, Hanson J, Safdi M, et al. Balsalazide is superior to mesalamine in the time to improvement of signs and symptoms of acute mild-to-moderate ulcerative colitis. *Am J Gastroenterol.* 2002; 97:3078-3086.

88 Pentasa [package insert]. Lexington, MA; Shire US; October 2015.

89 Pruitt R, Hanson J, Safdi M, et al. Balsalazide is superior to mesalamine in the time to improvement of signs and symptoms of acute mild-to-moderate ulcerative colitis. *Am J Gastroenterol.* 2002; 97:3078-3086.

90 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.

91 Kruis W, Judmaier G, Kayasseh L, et al. Double-blind dose-finding study of olsalazine versus sulphasalazine as maintenance therapy for ulcerative colitis. *Eur J Gastroenterol Hepatol.* 1995; 7:391-396.

92 Azulfidine EN-tabs [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.

93 Lialda [package insert]. Lexington, MA; Shire US; November 2015.

94 Asacol HD [package insert]. Rockaway, NJ; Warner Chilcott; October 2013.

95 Rowasa [package insert]. Somerset, NJ; Meda; June 2013.

96 sfRowasa [package insert]. Somerset, NJ; Meda; June 2013.

97 Canasa [package insert]. Bridgewater, NJ; Aptalis; December 2013.

98 Uceris [package insert]. Raleigh, NC; Salix; January 2015.

99 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.

100 Uceris [package insert]. Raleigh, NC; Salix; January 2015.

101 Gaffer MH, O'Brien CJ, Holdsworth CD. Clinical tolerance to three 5-aminosalicylic acid releasing preparations in patients with inflammatory bowel disease intolerant or allergic to sulphasalazine. *Aliment Pharmacol Ther.* 1992; 6:51-59.

102 Uceris [package insert]. Raleigh, NC; Salix; January 2015.

103 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.

104 Colazal [package insert]. Raleigh, NC; Salix; January 2012.

105 Giazio [package insert]. Raleigh, NC; Salix; December 2013.

106 Asacol [package insert]. Rockaway, NJ; Warner Chilcott; May 2015.

107 Delzicol [package insert]. Rockaway, NJ; Warner Chilcott; September 2015.

108 Pentasa [package insert]. Lexington, MA; Shire US; October 2015.

109 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.

110 Azulfidine [package insert]. New York, NY; Pharmacia & Upjohn; August.

111 Azulfidine EN-tabs [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.

112 Lialda [package insert]. Lexington, MA; Shire US; November 2015.

113 Apriso [package insert]. Morrisville, NC; Salix; July 2009.

114 Asacol HD [package insert]. Rockaway, NJ; Warner Chilcott; October 2013.

115 Rowasa [package insert]. Somerset, NJ; Meda; June 2013.

116 sfRowasa [package insert]. Somerset, NJ; Meda; June 2013.

-
- 117 Canasa [package insert]. Bridgewater, NJ; Aptalis; December 2013.
- 118 Uceris [package insert]. Raleigh, NC; Salix; January 2015.
- 119 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.
- 120 Ferry GD, Kirschner BS, Grand RJ, et al. Olsalazine versus sulfasalazine in mild to moderate childhood ulcerative colitis: results of the Pediatric Gastroenterology Collaborative Research Group Clinical Trial. *J Pediatr Gastroenterol Nutr.* 1993; 17(1):32-38.
- 121 Colazal [package insert]. Raleigh, NC; Salix; January 2012.
- 122 Giazio [package insert]. Raleigh, NC; Salix; December 2013.
- 123 Delzicol [package insert]. Rockaway, NJ; Warner Chilcott; September 2015.
- 124 Pentasa [package insert]. Lexington, MA; Shire US; October 2015.
- 125 Dipentum [package insert]. Somerset, NJ; Meda; January 2014.
- 126 Azulfidine [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.
- 127 Azulfidine EN-tabs [package insert]. New York, NY; Pharmacia & Upjohn; March 2014.
- 128 Lialda [package insert]. Lexington, MA; Shire US; November 2015.
- 129 Apriso [package insert]. Morrisville, NC; Salix; July 2009.
- 130 Asacol HD [package insert]. Rockaway, NJ; Warner Chilcott; October 2013.
- 131 sfRowasa [package insert]. Somerset, NJ; Meda; June 2013.
- 132 Rowasa [package insert]. Somerset, NJ; Meda; June 2013.
- 133 Canasa [package insert]. Birmingham, AL; Axcan Scandipharm; December 2013.
- 134 Uceris [package insert]. Raleigh, NC; Salix; January 2015.
- 135 Uceris [package insert]. Bridgewater, NJ; Salix; November 2015.
- 136 Delzicol [package insert]. Rockaway, NJ; Warner Chilcott; September 2015.
- 137 Lichtenstein GR, Gordon GL, Zakko S, et al. Clinical trial: once-daily mesalamine granules for maintenance of remission of ulcerative colitis – a 6-month placebo-controlled trial. *Aliment Pharmacol Ther.* 2010; 32(8): 990–999.
- 138 Sandborn WJ, Regula J, Feagan BG, et al. Delayed-release oral mesalamine 4.8 g/day (800 mg tablet) is effective for patients with moderately active ulcerative colitis. *Gastroenterology.* 2009; 137(6): 1934-43.
- 139 Green JR, Lobo AJ, Holdsworth CD, et al. Balsalazide is more effective and better tolerated than mesalamine in the treatment of acute ulcerative colitis. *Gastroenterology.* 1998; 114:15-22.
- 140 Levine DS, Riff DS, Pruitt R, et al. A randomized, double blind, dose-response comparison of balsalazide 6.75 g, balsalazide 2.25 g, and mesalamine 2.4 g in the treatment of active, mild-to-moderate ulcerative colitis. *Am J Gastroenterol.* 2002; 97:1398-1407.
- 141 Pruitt R, Hanson J, Safdi M, et al. Balsalazide is superior to mesalamine in the time to improvement of signs and symptoms of acute mild-to-moderate ulcerative colitis. *Am J Gastroenterol.* 2002; 97:3078-3086.
- 142 Procter & Gamble Pharmaceuticals. Data on file.
- 143 Giazio [package insert]. Raleigh, NC; Salix; December 2013.
- 144 Kruis W, Judmaier G, Kayasseh L, et al. Double-blind dose-finding study of olsalazine versus sulphasalazine as maintenance therapy for ulcerative colitis. *Eur J Gastroenterol Hepatol.* 1995; 7:391-396.
- 145 Kiillerich S, Ladefoged K, Rannem T, et al. Prophylactic effects of olsalazine v sulphasalazine during 12 months maintenance treatment of ulcerative colitis. The Danish Olsalazine Study Group. *Gut.* 1992; 33(2):252-255.
- 146 Lichtenstein GR, Kamm MA, Boddu P, et al. Effect of once or twice daily MMX mesalamine for the induction of remission of mild to moderately active ulcerative colitis. *Clin Gastroenterol Hepatol.* 2007; 5(1):95-102.
- 147 Kamm MA, Sandborn WJ, Gassull M, et al. Once-daily, high-concentration MMX mesalamine in active ulcerative colitis. *Gastroenterology.* 2007; 132(1):66-75.
- 148 D’Haens G, Sandborn WJ, Barrett K, et al. Once-daily MMX mesalamine for endoscopic maintenance of remission of ulcerative colitis. *Am J Gastroenterol.* 2012; 107(7):1064-77.
- 149 Uceris [package insert]. Raleigh, NC; Salix; January 2015.
- 150 Sandborn WJ, Travis S, Moro L, Jones R, et al. Once-daily budesonide MMX extended-release tablets induce remission in patients with mild to moderate ulcerative colitis: results from the CORE I study. *Gastroenterology.* 2012;143(5):1218.
- 151 Uceris [package insert]. Raleigh, NC; Salix; January 2015.
- 152 Uceris [package insert]. Bridgewater, NJ; Salix ; November 2015.
- 153 Sandborn WJ, Bosworth B, Zakko S, Gordon GL, et al. Budesonide induces remission in patients with mild to moderate ulcerative proctitis and ulcerative proctosigmoiditis. *Gastroenterology* 2015; 154 (15):00154-7. DOI: 10.1053/j.gastro.2015.01.037.
-